

2021

Transcendence of computer vision syndrome due to prolonged exposure to electronic devices

Klisman Custodio Sanchez

Facultad de Medicina Humana. Universidad Ricardo Palma, Lima-Perú,
lennon.custodio.sanchez@gmail.com

Follow this and additional works at: <https://inicib.urp.edu.pe/rfmh>

Recommended Citation

Custodio Sanchez, Klisman (2021) "Transcendence of computer vision syndrome due to prolonged exposure to electronic devices," *Revista de la Facultad de Medicina Humana: Vol. 21: Iss. 2, Article 29.*

DOI: <https://doi.org/10.25176/RFMH.v21i2.3611>

Available at: <https://inicib.urp.edu.pe/rfmh/vol21/iss2/29>

This Letter to the Editor is brought to you for free and open access by INICIB-URP. It has been accepted for inclusion in Revista de la Facultad de Medicina Humana by an authorized editor of INICIB-URP.



TRANSCENDENCE OF COMPUTER VISION SYNDROME DUE TO PROLONGED EXPOSURE TO ELECTRONIC DEVICES

TRANSCENDENCIA DEL SÍNDROME VISUAL INFORMÁTICO POR EXPOSICIÓN PROLONGADA A DISPOSITIVOS ELECTRÓNICOS

Custodio Sánchez K.^{1,a}

LETTERS TO THE EDITOR

Mr. Editor

Eye health, without a doubt, has been gaining more relevance in recent years due to the promotion of healthy habits in search of caring for our eyes, but the question also arises as to whether we are increasingly exposed to electronic devices with screens, these, can be detrimental to our visual well-being since, due to confinement, the use of these as a work and study tool has increased exponentially.

The American Optometric Association (AOA) defines Computer Visual Syndrome (CVS) or digital eyestrain as a "group of eye-related problems resulting from prolonged use of computers and cell phones". The term may not be completely accurate, but it is prevalent among computer users. The most important symptoms are: dry eyes, asthenopia, blurred vision, etc.⁽¹⁾.

It is estimated that the average American worker spends around seven hours a day in front of the computer (AOA) and 90% of the 70 million of these, who use computers for more than three hours a day, experience CVS in some form. Symptoms may be due to inadequate viewing distances, poor posture, or a combination of these factors⁽²⁾. The 2016 CVS report in the United States, where more than 10,000 adults participated, identified a prevalence of symptoms of 65%, with women being more affected than men (69% vs 60%, respectively). CVS was reported more frequently by people who used two or more devices simultaneously, with a 75% prevalence. The finding of greater symptoms in women was in line with the 2012 findings among a cohort of 520 office workers in New York and may be related to gender differences in the prevalence of CVS⁽³⁾.

In the European Union, the number of workers using information and communications technology (ICT) has been increasing⁽⁴⁾. The use of these devices is more widespread in the financial services sectors, who reported using it at a high intensity of up to 57%⁽⁴⁾.

Among 426 Spanish workers, the prevalence of CVS was 53%. With six or more hours of computer use, contact lens wearers were more likely to be affected than non-wearers, with prevalence of 65% and 50%, respectively. The finding was attributed to the mechanical interaction of the silicone hydrogel lenses with the ocular surface⁽³⁾.

The Peruvian context is no stranger to global data, there has been an increase in household access to ICT. In the second quarter of 2020, for every 100 households in 99 there is at least one ICT⁽⁵⁾. In relation to households with at least one computer in the first quarter of 2020, 94% is for exclusive household use⁽⁵⁾. There is not much specific information on CVS has been found in Peru, it may be possible to cite a study on the effect of exposure time to data display screens on visual fatigue in digitizers of the

¹ Facultad de Medicina Humana. Universidad Ricardo Palma, Lima-Perú.

^a Human Medicine student.

Cite as: Custodio Sánchez K. Transcendence of computer vision syndrome due to prolonged exposure to electronic devices. Rev. Fac. Med. Hum. April 2021; 21(2):463-464. DOI 10.25176/RFMH.v21i2.3611

Journal home page: <http://revistas.urp.edu.pe/index.php/RFMH>

Article published by the Magazine of the Faculty of Human Medicine of the Ricardo Palma University. It is an open access article, distributed under the terms of the Creative Commons License: Creative Commons Attribution 4.0 International, CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>), that allows non-commercial use, distribution and reproduction in any medium, provided that the original work is duly cited. For commercial use, please contact revista.medicina@urp.pe

Almenara Hospital in which it is concluded that there is a greater probability of presenting it due to the exposure time⁽⁶⁾. A second study, conducted at the Universidad Peruana Unión, shows the prevalence of CVS among all postgraduate university students (61%), due to computers (57.5%), followed by cell phone (37%)⁽⁷⁾.

Given that pedagogical and productive tasks require the use of ICT, it is possible to predict that CVS could

become a public health problem taking as a reference the world statistics that shows a high prevalence and adding the lack of knowledge of this problem, it will be of utmost importance to disseminate eye health awareness campaigns, as well as to educate the population on the use of adequate lighting levels, correct posture and periodic breaks, which will lead to an improvement in people's eye health.

Authorship contributions: The author participated in the conception, writing, final revision and approval of the manuscript.

Financing: Self-financed.

Interest conflict: The authors declare that they have no conflicts of interest in the publication of this article.

Received: October 18, 2020

Approved: January 02, 2021

Correspondence: Klisman Lennon Custodio Sánchez

Address: Av. Benavides 5440, Santiago de Surco, Lima 33 - Perú.

Telephone: (0511) 7080000

E-mail: klisman.custodio@urp.edu.pe

BIBLIOGRAPHIC REFERENCES

1. Ashish Chawla y cols. Computer Vision Syndrome: Darkness under the Shadow of Light. Canadian Association of Radiologists Journal 70 (2019) 5-9. doi.org/10.1016/j.carj.2018.10.005. <https://journals.sagepub.com/doi/10.1016/j.carj.2018.10.005>
2. Susan A. Randolph. Computer Vision Syndrome. Workplace Health Saf. 2017 Jul; 65(7):328. doi: 10.1177/2165079917712727. <https://pubmed.ncbi.nlm.nih.gov/28628753/>
3. Sheppard AL, Wolffsohn JS. Digital eye strain: prevalence, measurement and amelioration. BMJ Open Opth 2018;3:e000146. doi:10.1136/bmjophth-2018-000146. <https://pubmed.ncbi.nlm.nih.gov/29963645/>
4. Eurofound. Sixth European Working Conditions Survey. 2017. https://www.eurofound.europa.eu/sites/default/files/ef_publication/field_ef_document/ef1634en.pdf
5. INEI. Estadísticas de las tecnologías de información y comunicación en los hogares. INFORME TÉCNICO N° 03 – septiembre 2020. Instituto Nacional de Estadística e Informática - INEI. https://www.inei.gob.pe/media/MenuRecursivo/boletines/informe_tic_abr-may_jun2020.pdf
6. Vásquez García, Efecto del tiempo de exposición a pantallas de visualización de datos sobre la fatiga visual en digitadores del HNGAI -EsSALUD. UNIVERSIDAD NACIONAL MAYOR DE SAN MARCOS. FACULTAD DE MEDICINA. UNIDAD DE POST GRADO. <https://cybertesis.unmsm.edu.pe/handle/20.500.12672/2080>
7. Fernández Villacorta, Prevalencia del síndrome visual informático en estudiantes universitarios de postgrado de una universidad privada Lima -2019. UNIVERSIDAD PERUANA UNION. https://repositorio.upeu.edu.pe/bitstream/handle/UPEU/1153/Germ%C3%A1n_Tesis_Maestro_2017.pdf?sequence=5&isAllowed=y