

2021

The reality of patients in hemodialysis before COVID-19 pandemic

Mónica Vásquez Bambaren

Víctor Soto-Caceres

Victor Serna-Alarcón

Faculta de medicina humana Universidad Privada Antenor Orrego, Lima-Perú.,
victor.serna.alarcon@gmail.com

Follow this and additional works at: <https://inicib.urp.edu.pe/rfmh>

Recommended Citation

Vásquez Bambaren, Mónica; Soto-Caceres, Víctor; and Serna-Alarcón, Victor (2021) "The reality of patients in hemodialysis before COVID-19 pandemic," *Revista de la Facultad de Medicina Humana*: Vol. 21: Iss. 4, Article 27.

DOI: <https://doi.org/10.25176/RFMH.v21i4.4084>

Available at: <https://inicib.urp.edu.pe/rfmh/vol21/iss4/27>

This Letter to the Editor is brought to you for free and open access by INICIB-URP. It has been accepted for inclusion in Revista de la Facultad de Medicina Humana by an authorized editor of INICIB-URP.



THE REALITY OF PATIENTS IN HEMODIALYSIS BEFORE COVID 19 PANDEMIC

LA REALIDAD DEL PACIENTE EN HEMODIALISIS ANTES DE LA PANDEMIA COVID 19

Mónica Vásquez Bambaren^{1,a}, Víctor Soto-Caceres^{1,a,c}, Víctor Serna-Alarcón^{2,a,b}

Mr. Editor

In a pilot test carried out in the department of Lambayeque (Peru) about the profile of patients who entered hemodialysis for the first time with grade five chronic kidney disease (CKD 5), evidence was found different from other Latin American

The treatment modality for CKD 5 patients increased through hemodialysis according to the Latin American Registry of Dialysis and Kidney Transplantation from 2000 to 2008. In Peru, 84.75% of patients are registered with this form of treatment still in force in our country and others⁽¹⁾.

During admission to hemodialysis, the majority of patients are people with low socioeconomic status. In this population, health promotion and prevention is not intensified, being more susceptible to non-communicable chronic diseases⁽²⁾.

When these patients are admitted to hemodialysis for the first time, the clinical assessment is essential but epidemiological references must also be taken into account in order to have a better management. In laboratory tests, they are admitted with a hematocrit less than 30%, albumin less than 3.5 g / dl and with positive hepatitis C; the most common vascular access is venous catheter⁽³⁾. However, the pilot provides that the age, gender and vascular access coincide but the predominant causes are firstly arterial hypertension and secondly, diabetic nephropathy with a reason for admission of diabetic encephalopathy in the majority, being this last data where several studies do not take it into account; In the laboratory tests, a hematocrit of 23.53%, hemoglobin of 7.82, albumin of 3.12 gr / dl, creatinine of 11.17 and non-reactive to hepatitis C was found, reinforcing the information from the pilot, in a public hospital in our country in 2018, where it is obtained that in terms of sex it is slightly higher in men than in women, the age is less than 60 years, the causes are according to predominance as described above and in the case of hemoglobin it is less than 7g / dl⁽⁴⁾.

Due to the aforementioned, the doctor-patient relationship should be emphasized for a greater knowledge of associated comorbidities and possible complications that may occur during and after treatment; going hand in hand with mental health surveillance involving the family and caregiver for greater effectiveness in improving health in a holistic way; and in the same way, it takes into account the nutritional management in a diet that makes up for the deficit of proteins, vitamins and iron⁽⁵⁻⁶⁾.

In conclusion, the patient who is admitted to CKD 5 hemodialysis for the first time should be treated in an interdisciplinary manner through practical strategies by health personnel, especially by nephrologists, since this will allow, in the medium term, to be able to train the first level of care to thus avoid patients do not arrive in inadequate conditions to the emergency; and in the same way, information can be updated to keep track of reality and thus establish a health policy that helps to improve current management.

¹ Facultad de medicina humana Universidad San Martín de Porres, Lima-Perú.

² Facultad de medicina humana Universidad Privada Antenor Orrego, Lima-Perú.

^a Surgeon.

^b Master of Science in Clinical Research.

^c Doctor of Medicine.

Cite as: Mónica Vásquez Bambaren, Víctor Soto-Caceres, Víctor Serna-Alarcón. The reality of patients in hemodialysis before COVID 19 pandemic. Rev. Fac. Med. Hum. October 2021; 21(4):881-882. DOI 10.25176/RFMH.v21i4.4084

Authorship contributions: MVB, VSA have participated in the conception of the article, data collection, writing it, VRSC approved the final version in addition to contributing to the final writing.

Financing: Self-financed.

Conflict of interest: The authors declare that they have no conflict of interest with the manuscript and / or written content.

Received: August 15, 2021

Approved: September 04, 2021


Correspondence: Victor Serna Alarcón
Address: Mz C lote 28 Urb. Ingeniería, Trujillo.
Telephone: 969444641
Email: victor.serna.alarcon@gmail.com

BIBLIOGRAPHIC REFERENCES

1. Gonzalez-Bedat. Análisis del crecimiento de la prevalencia de las diferentes modalidades de tratamiento sustitutivo de la función renal en Latinoamérica. *Rev. Nefrol. Dial. Traspl.* 2014; 34 (4):170 –182.
2. García-García G, Jha V. Enfermedad renal crónica en poblaciones en situación de desventaja. *Rev. Nefrol. Dial. Traspl.* 2015; 35(1):1-6.
3. M. Marinovich. Características de la Población Incidente en Diálisis Crónica en Argentina. *Rev. Nefrol. Dial. Traspl.* 2011; 31(3): 104-117
4. Huauya-Leuyacc C, Palacios-Guillen AM, Benites-Zapata VA. Factores epidemiológicos asociados a la referencia temprana al nefrólogo para hemodiálisis crónica en pacientes de un Hospital Público de Perú. *Rev. Nefrol. Dial. Traspl.* 2018; 38(2):126-33.
5. Román D, Bustamante J. Aspectos nutricionales en la insuficiencia renal. *Nefrología.* 1 de julio de 2008; 28(3):333-42.
6. Candia, L. (2013). Seguimiento del paciente con enfermedad renal crónica en el primer nivel de atención de salud. Rosario de Santa Fe: Rev. Med. Rosario 2013, 79:28-38.


LETTERS TO THE EDITOR

Indexed in:




SciELO
Scientific Electronic Library Online

http://www.scielo.org.pe/scielo.php?script=sci_serial&pid=2308-0531&lng=es&nrm=iso




DIGITAL COMMONS

<https://network.bepress.com/>




DOAJ
DIRECTORY OF OPEN ACCESS JOURNALS

<https://doaj.org/>



LILACS
biblioteca virtual em saúde

<http://lilacs.bvsalud.org/es/2017/07/10/revistas-indizadas-en-lilacs/>



Revista de la Facultad de Medicina Humana

INSTITUTO DE INVESTIGACIÓN EN CIENCIAS BIOMÉDICAS

ISSN 2308-0531
DOI: 10.25176/RFMH.v21i4.4084