

2024

Short and prolonged fasting prior to the performance of tracheostomies in intensive therapy: a retrospective study

Lucas Gonzalo Duran
UNIVERSIDAD NACIONAL DEL SUR


María Emilia Beilman
UNIVERSIDAD NACIONAL DEL SUR

Araceli Natali Quiroga
HOSPITAL MUNICIPAL DE AGUDOS DR. LEONIDAS LUCERO

Magdalena Cruz
HOSPITAL MUNICIPAL DE AGUDOS DR. LEONIDAS LUCERO

Alejandra Vanesa Millan
HOSPITAL MUNICIPAL DE AGUDOS DR. LEONIDAS LUCERO

Follow this and additional works at: <https://inicib.urp.edu.pe/rfmh>

 [next page for additional authors](#)
Part of the [Health Information Technology Commons](#), [Medical Specialties Commons](#), and the [Public Health Commons](#)

Recommended Citation

Gonzalo Duran, Lucas; Emilia Beilman, María; Natali Quiroga, Araceli; Cruz, Magdalena; Vanesa Millan, Alejandra; Johanna Ojeda, Micaela; Ciccioli, Fabiana; Montenegro Fernandez, Micaela Giselle; Monrroy Miro, Wendy Estefany; Trinidad Malisia, Valentina; Antonio Grassi, Nicolas; Zelaya de Leon, Nazareno Iñaki; Ezequiel Espinoza, Franco; Otamendi, Marina; Zorzano Osinalde, Paula; and Petasny, Marcos (2024) "Short and prolonged fasting prior to the performance of tracheostomies in intensive therapy: a retrospective study," *Revista de la Facultad de Medicina Humana*: Vol. 24: Iss. 1, Article 7.
DOI: <https://doi.org/10.25176/RFMH.v24i1.6086>
Available at: <https://inicib.urp.edu.pe/rfmh/vol24/iss1/7>

This Article is brought to you for free and open access by INICIB-URP. It has been accepted for inclusion in Revista de la Facultad de Medicina Humana by an authorized editor of INICIB-URP.

Short and prolonged fasting prior to the performance of tracheostomies in intensive therapy: a retrospective study

Authors

Lucas Gonzalo Duran, María Emilia Beilman, Araceli Natali Quiroga, Magdalena Cruz, Alejandra Vanesa Millan, Micaela Johanna Ojeda, Fabiana Ciccicoli, Micaela Giselle Montenegro Fernandez, Wendy Estefany Monroy Miro, Valentina Trinidad Malisia, Nicolas Antonio Grassi, Nazareno Iñaki Zelaya de Leon, Franco Ezequiel Espinoza, Marina Otamendi, Paula Zorzano Osinalde, and Marcos Petasny



SHORT AND PROLONGED FASTING PRIOR TO THE PERFORMANCE OF TRACHEOSTOMIES IN INTENSIVE THERAPY: A RETROSPECTIVE STUDY

AYUNO CORTO VS PROLONGADO PREVIO A LA REALIZACION DE TRAQUEOSTOMÍAS EN PACIENTES DE TERAPIA INTENSIVA: UN ESTUDIO RETROSPECTIVO

Lucas Gonzalo Duran ^{1,2}, María Emilia Beilman ^{1,2}, Araceli Natali Quiroga ^{1,3}, Magdalena Cruz ^{1,2}, Alejandra Vanesa Millan ^{1,2}, Micaela Johanna Ojeda ^{1,2}, Fabiana Ciccioli ^{1,2}, Micaela Giselle Montenegro Fernandez ^{1,2}, Wendy Estefany Monrroy Miro ^{1,2}, Valentina Trinidad Malisia ^{1,2}, Nicolas Antonio Grassi ^{1,2}, Nazareno Iñaki Zelaya De Leon ^{1,2}, Franco Ezequiel Espinoza ^{1,2}, Marina Otamendi ^{1,2}, Paula Zorzano Osinalde ^{1,2}, Marcos Petasny ^{1,2}

ABSTRACT

Introduction: International guidelines on aid prior to invasive procedures usually generate longer aid than in intensive care (IT) patients. This fact represents a high risk of malnutrition and, consequently, a worse prognosis. The objective of the present investigation was to analyze the degree of association between the fasting time prior to tracheostomy of patients under mechanical ventilatory assistance (MVA) and the appearance of pneumonia. **Methods:** Retrospective cohort study that included patients admitted to our IT from 10/01/2018 to 08/31/2022 and with a tracheostomy performed. Two cohorts were defined characterized by fasting ≤ 3 hours vs. > 3 hours. Fisher's exact test and Mann-Whitney test were used for bivariate analysis. A p value < 0.05 was shown to be significant. **Results:** 141 patients were hospitalized with a tracheostomy, 9 were excluded, leaving 132 patients. The cohort with fasting ≤ 3 hours was made up of 15 patients and the one with fasting > 3 hours was made up of 117, the latter presented an average fast of 2.5 hours (IR 2-3), days of AVM prior to the procedure of 13 days (IR 12-18), while the other cohort presented an average fast of 6 hours (IR 5-8), days of AVM prior to the procedure of 12 days (IR 10-14.5). When analyzing the association between the type of fasting and the appearance of pneumonia, an OR of 0.958 (95% CI: 0.32-2.87) was obtained, p value of 0.743. **Conclusions:** No significant differences were found regarding fasting time and the appearance of pneumonia as referred to in the international literature.

Keywords: Fasting, Tracheostomy, Pneumonia. (Source: MESH-NLM)

RESUMEN

Introducción: Las pautas internacionales sobre la práctica del ayuno previo a procedimientos invasivos generalmente ocasionan ayunos más prolongados que en los casos de pacientes internados en las Unidades de Cuidados Intensivos (UCI). Este hecho representa un alto riesgo de desnutrición y, consecuentemente, un pronóstico más negativo. El objetivo de la presente investigación fue analizar el grado de asociación entre el tiempo de ayuno previo a la traqueostomía de pacientes bajo asistencia ventilatoria mecánica (AVM) y la aparición subsecuente de neumonía. **Métodos:** Fue un estudio de cohorte retrospectivo que incluyó a pacientes ingresados en nuestra UCI desde el 01/10/2018 hasta el 31/08/2022, quienes habían sido sometidos a una traqueostomía. Se definieron dos cohortes caracterizadas por ayuno inferior o igual a tres horas y superior a tres horas. Se utilizó la prueba exacta de Fisher y la prueba U de Mann-Whitney para el análisis bivariado. Un valor de $p < 0.05$ se consideró significativo. **Resultados:** Se hospitalizaron 141 pacientes con traqueostomía, 9 fueron excluidos, quedando 132 pacientes. La cohorte con ayuno ≤ 3 horas estuvo compuesta por 15 pacientes y la de ayuno > 3 horas estuvo compuesta por 117; esta última presentó un ayuno promedio de 2.5 horas (RIC 2-3) y 13 días de AVM previos al procedimiento (RIC 12-18), mientras que la otra cohorte presentó un ayuno promedio de 6 horas (RIC 5-8) y 12 días de AVM previos al procedimiento (RIC 10-14.5). Al analizar la asociación entre el tipo de ayuno y la aparición de neumonía, se obtuvo un OR de 0.958 (IC del 95%: 0.32-2.87) y valor de p de 0.743. **Conclusiones:** No se encontraron diferencias significativas respecto al tiempo de ayuno y la aparición de neumonía, tal como se refleja en la literatura internacional.

Palabras clave: Ayuno; Traqueostomía; Neumonía. (Fuente: DeCS- BIREME)

¹ Departamento de Ciencias de la Salud, Universidad Nacional del Sur, Bahía Blanca, Buenos Aires, Argentina.

² Servicio de Terapia intensiva, Hospital municipal de agudos Dr. Leonidas Lucero, Bahía Blanca, Buenos Aires, Argentina.

³ Servicio de cirugía, Hospital municipal de agudos Dr. Leonidas Lucero, Bahía Blanca, Buenos Aires, Argentina.

Cite as : Duran Lucas G, Beilman ME, Quiroga AN, Cruz M, Millan AV, Ojeda MJ, Ciccioli F, Montenegro Fernandez MG, Monrroy Miro WE, Malisia VT, Grassi NA, Zelaya De Leon NI, Espinoza FE, Otamendi M, Zorzano Osinalde P, Petasny M. Short and Prolonged fasting prior to the performance of Tracheostomies in intensive therapy: a retrospective study. Rev Fac Med Hum. 2024;24(1):58-64. doi 10.25176/RFMH.v24i1.6086

Journal home page: <http://revistas.urp.edu.pe/index.php/RFMH>

Article published by the Journal of the Faculty of Human Medicine of the Ricardo Palma University. It is an open access article, distributed under the terms of the Creative Commons License: Creative Commons Attribution 4.0 International, CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>), which allows non-commercial use, distribution and reproduction in any medium, provided that the original work is duly cited. For commercial use, please contact revista.medicina@urp.edu.pe





INTRODUCTION

Tracheostomy is one of the most frequently performed procedures in the intensive care unit (ICU). According to medical literature, it is performed in 9-10 % of all mechanically ventilated patients in the United States and the United Kingdom, respectively^(1,2,3,4). However, other studies that evaluated its international prevalence have reported higher values, arguing that 24 % of these patients were assisted through a tracheostomy⁽⁵⁾.

Generally, these patients, who are being evaluated to undergo a tracheostomy, are usually in an extremely critical condition with multiple associated comorbidities. This is aggravated when the procedures are delayed due to the lack of fasting time of more than eight hours, indicated by the anaesthesiologists or surgeons, that causes patients to remain fasting for more than a day in many cases, with a high metabolic demand and, therefore, a greater risk of malnutrition, a condition that increases the risk of septic complications and insulin resistance syndrome^(6,7). This fact produces a delay in transfers to less complex floors and prolongs stays in the ICU, which is associated to an increase in care costs.

Contrary to what may be assumed, pulmonary aspiration during anesthesia is a very rare event with low incidence^(8,9). Bronchoaspiration is an unusual adverse circumstance, with an incidence of 2.9 to 4.7 cases per 10 000 general anesthesia in the overall population⁽¹⁰⁾. Due to the physiological changes that occur during pregnancy, obstetric patients have a two to three times higher risk⁽¹¹⁾. Bronchoaspiration can be classified into two categories: aspiration pneumonia, which refers to undetected aspiration of gastrointestinal contents that triggers pneumonia, and aspiration pneumonitis, which refers to aspiration of gastric contents, that is directly observed by the doctor during anesthetic induction, the surgical procedure or immediately after it⁽¹²⁾. When aspiration occurs, it is most likely related to a complication during induction of anesthesia, laryngoscopy, or extubation⁽¹³⁾. For this reason, it is possible to consider shorter preoperative fasting times for patients, who are already endotracheally intubated and have easy access to the

stomach through nasogastric or orogastric tubes. The aim of the present investigation is to analyze the degree of association between the fasting time prior to tracheostomy of patients under mechanical ventilatory assistance (MVA) and the subsequent appearance of pneumonia, mortality from all causes, length of hospital stays, duration of intensive care stays and the decline of the PAFI.

METHODS

A retrospective cohort study was performed. The target population was made up of all patients hospitalized in a period between October 1, 2018 and August 31, 2022 in ICU of the Hospital Municipal de Agudos Dr. Leónidas Lucero (HMALL) de Bahía Blanca.

Unit of analysis or eligible population

Inclusion criteria: Only patients, who during their hospitalization, had a tracheostomy performed with any technique during the period in the study were included.

Exclusion criteria

Patients who required the emergency procedure were excluded.

Elimination criteria

Patients who, due to insufficient data recorded in the electronic medical record, were unable to obtain at least 80 % of the required data were eliminated.

Data collection instrument

Data were collected through the HMALL computer system that provides access to the digital medical history of each patient. Likewise, laboratory data were obtained through the use of the NextLab® computer system with which the hospital has an agreement.

Operational definition of the variables

Reason for hospitalization: in ICU and was confirmed in the epicrisis upon discharge. It was classified into: cardiological, neurological, respiratory, infectious and other clinical emergency, planned post-operative, non-neurological surgical emergency and post-surgical complications.



Type of fast: fasts were classified as short if they were less or equal than three hours and prolonged when they exceeded that period.

Associated pneumonia: defined by suggestive symptoms (fever, purulent mucus, crackles), decrease in PaFi <200 , and chest x-ray with clear infiltrate. To capture all intervention-related pneumonias, the criteria were expanded to include any new consolidation within a 5-day period after the intervention.

PAFI: estimated by the ratio between the arterial oxygen pressure and the inspired fraction of oxygen. They were classified as pafi with severe decrease at values <100 , moderate between 100 and 200 and mild between 200 and 300.

MVA time: established in days until weaning from the ventilator or death.

Hospitalization time in ICU: established as the duration in days of the patients' hospitalization in the ICU.

Hospital stay time: established as the duration in days of the patients' stay in the hospital.

Reason for orotracheal intubation: This variable is defined as the reason why it was decided to protect the airway. This may be due to respiratory failure, ventilatory failure or cardiorespiratory arrest.

Type of tracheostomy: tracheostomy was classified if it was performed before 10 days and late if it was performed after that period.

Tracheostomy technique: they can be defined as open or percutaneous.

Surgical team: It could be composed of a resident – resident team; staff – resident; staff – staff.

Type of surgical wound: classified as clean wound, clean contaminated wound, contaminated wound and dirty wound.

Duration of the procedure: quantified in minutes from the beginning of the patient's induction until the fixation of the tracheostomy tube with suture.

Statistic analysis

Patients were coded numerically consecutively in consistent order with their medical record number, which guaranteed the anonymity and confidentiality of the information at the time of analysis. A univariate analysis was performed to estimate the percentages and means of each variable. To compare the means of the baseline characteristics, the Student T test was used, after verifying the assumptions for the delta PaFi variables, the rest of the numerical variables that did not have a normal distribution were analyzed with the Mann-Whitney U test. To estimate the degree of association between fasting time and pneumonia, a bivariate analysis was performed with cross-classification tables using Fisher's exact test and subsequently a logistic regression model was performed to analyze the factors associated with the presence of post-fasting pneumonia tracheostomy. A p value <0.05 was considered significant and the IBM SPSS 23® software in its version for Windows® was used.

Ethics statement

This research was approved by the Institutional Research Bioethics Committee of the HMALL (resolution No. 6-263-2022), accredited by the Central Ethics Committee of the Ministry of Health of the provincia de Buenos Aires, under No. 105/2013, with last reaccreditation on 09/09/2022. Strict confidentiality was guaranteed in the handling of the data provided by the personal data protection law 25 326 and was conducted in accordance with the guidelines established by the modified declaration of Helsinki.

RESULTS

During the time period covered by the study, a total of 1106 patients were admitted to the ICU, of which 12.75 % (141/1106) required a tracheostomy. Of these 141 patients, 9 were excluded and those are a total of 132. Two cohorts were established based on fasting time (>3 hours or ≤ 3 hours). The cohort with fasting ≤ 3 hours was made up of 15 patients, while the cohort with fasting >3 hours was made up of 117.





The cohort with fasting ≤ 3 hours had a median age of 53 years (IR 16), a proportion of male patients of 80 % (12/15), a median number of days of MVA until the procedure of 13 days (IR 6), a median fasting time of 2.5 hours (IR 1) and a positive previous pneumonia diagnosis of 25 % (3/15). Regarding the reason for hospitalization, it was identified that within this group 40% (6/15) presented a respiratory emergency, 33.3 % (5/15) a neurological emergency, and 13.3 % (2/15) a non-neurological surgical emergency, 6.7% (1/15) had a cardiological emergency and 6.7 % (1/15) had other clinical emergencies, while the most frequent reason for intubation was respiratory failure with 60 % (9/15), followed by ventilatory failure with 33.3 % (5/15) and cardio-respiratory arrest with 6.7 % (1/15).

On the other hand, the cohort with fasting > 3 hours had a median age of 61 years (IR 27.5), a proportion of male patients of 59.8 % (70/117), a median number of days of MVA until the procedure of 12 days (IR 5), a median fasting time of 6 hours (IR 4) and a previous positive pneumonia diagnosis of 35.89 % (42/117). Regarding

the reason for hospitalization, it was identified that within this group 26.5 % (31/117) presented a respiratory emergency, 31.6 % (37/117) a neurological emergency, and 14.5 % (17/117) a non-neurological surgical emergency, 7.7 % (9/117) had a cardiological emergency and 9.4 % (11/117) had other clinical emergencies, while the most frequent reason for intubation was respiratory failure with 52.1 % (61/117), followed by ventilatory failure with 42.7 % (50/117) and cardio-respiratory arrest with 5.1 % (6/117). When comparing the baseline characteristics previously described in each cohort, p values > 0.05 were obtained for the statistical tests used in each of the variables.

Regarding the bivariate analysis, when analyzing the association between the appearance of post-tracheostomy pneumonia according to the fasting time, an OR of 1.043 (95% CI: 0.348-3.124) was obtained with a P value for the chi-square test of 0.585. The analysis of the rest of the variables of interest can be seen summarized in table 1 and table 2.

Tabla 1. Comparative analysis for quantitative variables.

Characteristics	Fasting ≤ 3 hs	Fasting > 3 hs	Total	P value*
Total IT hospitalization days	25 (21-31)	25 (19.5-34)	25 (20-33)	0.943
Days of hospital stay	33 (25-49)	33 (25-52)	33 (25-50)	0.810
Days of AVM Totals	22 (19-29)	23 (18-32.5)	23 (18.25-31)	0.980
PAFI 24 hours after tracheostomy	232 (174-292)	208 (173.5-258.3)	212 (174-260)	0.296
PAFI 48 hours after tracheostomy	198 (179-280)	206 (170-263)	206 (171.25-264.75)	0.871
PAFI 72 hours after tracheostomy	235 (180-290)	220 (184-302)	223 (183-297.25)	0.949
Procedure duration (in minutes)	30 (20-40)	30 (20-30)	28,56 (20-30)	0.156

*U de Mann-Whitney

Tabla 2. Analysis of association between the different variables and the type of fasting.

Variable	Fasting ≤ 3 hs	Fasting > 3 hs	OR	P value*
No post-tracheostomy pneumonia	9	69	1.043 (0.348-3.124)	0.585
With post tracheostomy pneumonia	6	48		
Early tracheostomy	2	38	0.32 (0.069-1.489)	0.129
Late tracheostomy	13	79		
Does not die	9	59	1.475 (0.493-4.407)	0.485
Died	6	58		
Negative x-ray	6	59	0.655 (0.219-1.958)	0.447



Positive x-ray	9	58	0.655 (0.219-1.958)	0.447
Fever	11	83		
Febrile	4	34	1.127 (0.335-3.785)	0.847
Negative tracheal aspiration	12	95		
Positive tracheal aspiration	3	22	0.926 (0.241-3.564)	0.911
Surgical team: Resident - Resident	3	19		
Surgical team: Staff - Resident	9	73		0.934
Surgical team: Staff - Staff	3	25		
Clean wound	0	9		
Clean contaminated wound	12	86		0.671
Contaminated wound	1	11		
Dirty wound	2	11		

* Fisher Test

Given the retrospective characteristic of the established cohorts, a power of 12.6 % was calculated for the pooled sample. The multivariate analysis multiple regressions showed the same trend and obtained an OR of 0.90 (95 % CI: 0.69; 1.17), 0.81 (95 % CI: 0.40; 1.64) and 2.19 (95 % CI: 0.97; 4.96) for the reason for hospitalization, reason for intubation and previous pneumonia.

DISCUSSION

This study represents a valuable approach when examining the impact of fasting in patients with tracheostomies, given that there is little scientific evidence available on this topic. Although fasting before tracheostomy is a common practice in daily clinical care, there is a lack of solid local evidence that justifies the current fasting protocol, making it crucial to consider this aspect as a relevant indicator when evaluating the quality of tracheostomy medical attention. The results of this study highlight the high frequency of patients requiring tracheostomy in our service, coinciding with international data where references are made to similar statistics describing a 10 to 14 % frequency of this procedure (1,1).

The results of our study provide relevant evidence about the safety of performing the with a short fast without increasing the risk of developing pneumonia an prolongs the ICU or hospital stay or worsening mortality. These data are consistent with what was expressed in the experience published by Hartl T et al.,

who implemented a protocol without fasting prior to performing the procedure and found results clearly in favor of the alternative mentioned before⁽¹⁾. It should be noted that currently there is no clear indication of the most appropriate time to avoid complications and guarantee a good nutritional intake in these patients, which is supported by evidence. In fact, in Gonik's study carried out in 2016⁽¹⁾; statistically significant differences have not been found when comparing the group that received a 40-minute fast compared to the control group that received a 6-hour fast.

On the other hand, a survey conducted in the critical care and anesthesia departments found significant variability in internal protocols. Surgical ICU were noted to have shorter fasting periods for procedures compared to ICU and anesthesia departments (4 vs. 6 hours). However, in all specialties, wide fasting ranges were observed for tracheostomies (between 0-8 hours)⁽¹⁾. Likewise, anesthesia guidelines have few references to fasting times, especially in critically ill patients. Shime et al. found that 90 % of Japanese anesthesia departments used longer fasts than recommended⁽²¹⁾. Others have noted that fasts for solids actually ranged from 12 to 14 hours instead of 6 to 8 hours and can last up to 37 hours^(20,21).

Pousman et al. described short fasts for patients receiving jejunal feedings⁽²²⁾. Using a 45-minute fast before bedside procedures, they noted no change in





ventilator-associated pneumonias compared to controls and a trend toward increased caloric intake. In a review of aspiration, Warner et al identified only 68 cases of aspiration in 215 000 general anesthesia procedures⁽¹³⁾. All aspirations occurred before, after, or during airway manipulations and during induction or emergence of anesthesia. At such times, the airway is more susceptible to aspiration of gastric contents, since the patient is lying down with abolished protective reflexes and low esophageal tone. Even in high-risk emergency intubations, the factor that predisposes patients to the highest risk of aspiration is multiple intubations attempts and the need for mask ventilation that can inflate the stomach⁽²³⁾.

These promising results should be considered the first step towards a change of approach in the ICU. It is essential to generate solid evidence before implementing any change in clinical practice, so it is planned to continue this study with a prospective analysis, taking into account the limitations identified in this work. One of the limitations of our study lies in its retrospective design, so we recognize the need to develop a protocol that allows prospective follow-up of patients, which will surely improve the quality and

reliability of the data in question, adding nutritional parameters that enrich the debate. This prospective approach will allow us to collect more detailed and precise information on the effects of fasting in patients with tracheostomies and improve the sample size necessary to increase the power of the study given the low incidence of events.

CONCLUSIONS

No significant differences were found regarding the risk of pneumonia, days of hospital stay, PAFI values or mortality with respect to fasting time prior to the tracheostomy procedure.

It is important to highlight the importance of nutrition in critically ill patients, so it is pertinent to optimize fasting times. This is consistent with the limited results obtained in other countries, which adapted to our context and allows us to avoid postponing the procedure.

Data availability

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Authorship contribution: Conceptualization: DLG, BME, and QAN. Acquisition, analysis, or interpretation of the data: DLG, BME, QAN. Statistical analysis: DLG and OMJ. Drafting of the manuscript and editing: DLG, BME, QAN, CM, MAV, OMJ, CF, MVT, MFMG, MMWE, GNA, ZLNI, EFE, OM, ZOP, PM.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Received: December 04, 2023

Approved: February 28, 2024

Correspondence: Lucas Gonzalo Duran.

Address: Estomba 968, Bahía Blanca (8000), Buenos Aires, Argentina.

E-mail: lucas.duran@uns.edu.ar

REFERENCES

- McGrath BA, Bates L, Atkinson D, Moore JA. National Tracheostomy Safety Project. Multidisciplinary guidelines for the management of tracheostomy and laryngectomy airway emergencies. *Anaesthesia*. 2012;67:1025–41. DOI: [10.1111/j.1365-2044.2012.07217.x](https://doi.org/10.1111/j.1365-2044.2012.07217.x)
- Veenith T, Ganeshamoorthy S, Standley T, Carter J, Young P. Intensive care unit tracheostomy: A snapshot of UK practice. *Int Arch Med*. 2008;1:21. DOI: [10.1186/1755-7682-1-21](https://doi.org/10.1186/1755-7682-1-21)
- Krishnan K, Elliot SC, Mallick A. The current practice of tracheostomy in the United Kingdom: A postal survey. *Anaesthesia*. 2005;60:360–4. DOI: [10.1111/j.1365-2044.2004.04106.x](https://doi.org/10.1111/j.1365-2044.2004.04106.x)
- Mehta AB et al. Trends in tracheostomy for mechanically ventilated patients in the United States, 1993–2012. *Am J Respir Crit Care Med*. 2015;192:446–54. doi: [10.1164/rccm.201502-0239OC](https://doi.org/10.1164/rccm.201502-0239OC)
- Abe T, Madotto F, Pham T et al. Epidemiology and patterns of tracheostomy practice in patients with acute respiratory distress syndrome in ICUs across 50 countries. *Crit Care* 22, 195 (2018). <https://doi.org/10.1186/s13054-018-2126-6>.
- Tanaka A, Uchiyama A, Kitamura T et al. Association between early tracheostomy and patient outcomes in critically ill patients on mechanical ventilation: a multicenter cohort study. *J Intensive Care* 10, 19 (2022). <https://doi.org/10.1186/s40560-022-00610-x>



7. Hartl T, Anderson D, Levi J. Safety of a no-fast protocol for tracheotomy in critical care. *Can J Surg.* 2015 Feb;58(1):69-70. <https://doi.org/10.1503/cjs.027213>
8. Gonik N, Tassler A, Ow TJ, Smith RV, Shuaib S, Cohen HW, Sarta C, Schiff BA. Randomized Controlled Trial Assessing the Feasibility of Shortened Fasts in Intubated ICU Patients Undergoing Tracheotomy. *Otolaryngol Head Neck Surg.* 2016 Jan;154(1):87-93. <https://doi.org/10.1177/0194599815611859>
9. De Luca M y col. Duración del ayuno preoperatorio en pacientes con cirugía programada. *Rev. Hosp. Ital. B.Aires* 2019;39(3):77-80.
10. Ng A, Smith G. Gastroesophageal reflux and aspiration of gastric contents in anesthetic practice. *Anesth Analg.* 93 (2001), pp. 494-513. DOI: [10.1097/00000539-200108000-00050](https://doi.org/10.1097/00000539-200108000-00050)
11. Hayashi R, Maeda S, Hideki T, Higuchi H, Miyawaki T. Pulmonary Aspiration During Induction of General Anesthesia. *Anesth Prog.* 2020 Dec 1;67(4):214-218. doi: [10.2344/anpr-67-02-03](https://doi.org/10.2344/anpr-67-02-03)
12. Beck-Schimmer B, Bonvini JM. Bronchoaspiration: Incidence, consequences and management. *Eur J Anaesthesiol.* 28 (2011), pp. 78-84. DOI: [10.1097/EJA.0b013e32834205a8](https://doi.org/10.1097/EJA.0b013e32834205a8)
13. Sun J, Wei G, Hu L, Liu C, Ding Z. Perioperative pulmonary aspiration and regurgitation without aspiration in adults: a retrospective observational study of 166,491 anesthesia records. *Ann Palliat Med* 2021;10(4):4037-4046. <https://doi.org/10.21037/apm-20-2382>
14. Abe T, Madotto F, Pham T et al. Epidemiology and patterns of tracheostomy practice in patients with acute respiratory distress syndrome in ICUs across 50 countries. *Crit Care* 22, 195 (2018). <https://doi.org/10.1186/s13054-018-2126-6>.
15. Tanaka A, Uchiyama A, Kitamura T et al. Association between early tracheostomy and patient outcomes in critically ill patients on mechanical ventilation: a multicenter cohort study. *J Intensive Care* 10, 19 (2022). <https://doi.org/10.1186/s40560-022-00610-x>
16. Hartl T, Anderson D, Levi J. Safety of a no-fast protocol for tracheotomy in critical care. *Can J Surg.* 2015 Feb;58(1):69-70. <https://doi.org/10.1503/cjs.027213>
17. Gonik N, Tassler A, Ow TJ, Smith RV, Shuaib S, Cohen HW, Sarta C, Schiff BA. Randomized Controlled Trial Assessing the Feasibility of Shortened Fasts in Intubated ICU Patients Undergoing Tracheotomy. *Otolaryngol Head Neck Surg.* 2016 Jan;154(1):87-93. <https://doi.org/10.1177/0194599815611859>
18. De Luca M y et al.. Duración del ayuno preoperatorio en pacientes con cirugía programada. *Rev. Hosp. Ital. B.Aires* 2019;39(3):77-80.
19. Shime N, Ono A, Chihara E et al. Current practice of preoperative fasting: a nationwide survey in Japanese anesthesia-teaching hospitals. *J Anesth* 19, 187-192 (2005). <https://doi.org/10.1007/s00540-005-0319-z>
20. Anderson M, Comrie R. Adopting preoperative fasting guidelines. *AORN J.* 2009 Jul;90(1):73-80. <https://doi.org/10.1016/j.aorn.2009.01.026>
21. Crenshaw JT, Winslow EH. Preoperative fasting: old habits die hard. *Am J Nurs.* 2002 May;102(5):36-44; quiz 45. <https://doi.org/10.1097/00000446-200205000-00033>
22. Pousman RM, Pepper C, Pandharipande P, Ayers GD, Mills B, Diaz J, Collier B, Miller R, Jensen G. Feasibility of implementing a reduced fasting protocol for critically ill trauma patients undergoing operative and nonoperative procedures. *JPEN J Parenter Enteral Nutr.* 2009 Mar-Apr;33(2):176-80. <https://doi.org/10.1177/0148607108327527>
23. Sakles JC, Chiu S, Mosier J, Walker C, Stolz U. The importance of first pass success when performing orotracheal intubation in the emergency department. *Acad Emerg Med.* 2013 Jan;20(1):71-8. <https://doi.org/10.1111/acem.12055>

